



Negative Air Ions and Their Effects on Human Health and Air Quality Improvement

[Shu-Ye Jiang](#), [Ali Ma](#), and [Srinivasan Ramachandran](#)*

Temasek Life Sciences Laboratory, 1 Research Link, National University of Singapore, Singapore 117604, Singapore; shuye@tll.org.sg (S.-Y.J.); zhigang@tll.org.sg (A.M.)

* Correspondence: sri@tll.org.sg; Tel.: +65-6872-7480

Received 2018 Aug 15; Accepted 2018 Sep 25.

[Copyright](#) © 2018 by the authors.

Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Abstract

Negative air ions (NAIs) have been discovered for more than 100 years and are widely used for air cleaning. Here, we have carried out a comprehensive reviewing on the effects of NAIs on humans/animals, and microorganisms, and plant development. The presence of NAIs is credited for increasing psychological health, productivity, and overall well-being but without consistent or reliable evidence in therapeutic effects and with controversy in anti-microorganisms. Reports also showed that NAIs could help people in relieving symptoms of allergies to dust, mold spores, and other allergens. Particulate matter (PM) is a major air pollutant that affects human health. Experimental data showed that NAIs could be used to high-efficiently remove PM. Finally, we have reviewed the plant-based NAI release system under the pulsed electric field (PEF) stimulation. This is a new NAI generation system which releases a huge amount of NAIs under the PEF treatment. The system may be used to freshen indoor air and reduce PM concentration in addition to enriching oxygen content and indoor decoration at home, school, hospital, airport, and other indoor areas.

Keywords: negative air ions, superoxide, particulate matter, pulsed electric field

1. Introduction

Negative air ions (NAIs) have been discovered for more than 100 years [[1](#)]. Now, NAI generators are widely available for home or industrial uses. In the meantime, various new technologies were developed and used to further improve NAI generation and reduce the release of its byproduct ozone. However, some controversial results or comments have been reported for the beneficial effect on humans/animals or on reductions in bacterial densities. Here, we have carried out a comprehensive reviewing on NAIs. On the other hand, strong evidence had shown the roles of NAIs in high-efficiently reducing particulate matter (PM) concentration. Thus, more work should be done to further improve NAI release by new methods or devices so that NAIs could be more widely used for air cleaning. Here, we review the generation of NAIs and their effects on humans, animals, and microorganisms. We then discussed the involvement of superoxide ions in biological effects of NAIs. Subsequently, we focused

on plant-based NAI generation systems as these are relatively new NAI generation systems with some advantages on traditional corona discharge NAI generators. We also reviewed the air cleaning ability of NAIs, especially in removing PM with diameters less than 10 micrometers (PM₁₀).

2. Systematic Review of Literatures

The systematic review of studies on NAIs initiated by literature searches. Three databases were selected including PubMed database (available online: <https://www.ncbi.nlm.nih.gov/pubmed/advanced>), ScienceDirect database (available online: <https://www.sciencedirect.com/search/advanced>), and IEEE Xplore Digital Library (available online: <https://ieeexplore.ieee.org/search/advsearch.jsp>). The coverage period of the database search is around 100 years from 1918 to 2018 (the updated search date is 10 September 2018). Three keywords “negative”, “air”, and “ion” were used to search all collected articles in these databases. The database searches harvested a total of 335, 681, and 221 articles that contain all of the three keywords in either titles or abstracts when the PubMed database, ScienceDirect, and IEEE Xplore databases were employed, respectively. Only peer-reviewed and English-language articles were considered. We then screened these articles by manually reviewing titles, which led to 170, 279, and 117 studies selected (duplicated references were excluded); the remaining were excluded due to no relationship to the NAI topic. We further carried out abstract screening to exclude more unrelated articles. Based on the abstract reviewing, 93, 113, and 57 references were selected, which were subjected to full-length reviewing. During full-length article reviewing, some cited references, which were not included in the full-length review list, were also selected for additional reviewing.

3. Negative Air Ions and Their Generation

Air ions are electrically charged molecules or atoms in the atmosphere [2]. An air ion is formed when a gaseous molecule or atom receives sufficiently high energy to eject an electron [3]. NAIs are those that gain an electron, while positive air ions lose an electron. The natural and artificial energy sources include (1) radiant or cosmic rays in the atmosphere; (2) sunlight including ultraviolet; (3) natural and artificial corona discharge including thunder and lightning; (4) the shearing forces of water (Lenard effect); (5) plant-based sources of energy.

3.1. Radiant or Cosmic Rays in the Atmosphere

The radioactive elements such as uranium, radium, actinium, and thorium widely exist in our planet. They decay in the atmosphere and emit α , β , and/or γ rays, which ionize the air. Thus, radiant and cosmic ray ionization is ubiquitous in the Earth’s atmosphere. The cosmic ray ionization accounts for around 20% of the total ionization over land surfaces [4]. They are also the principal energy sources that generate NAIs over the oceans [5]. The concentration of NAIs produced by these rays might reach from 500 ions per cm³ in land surface [6] to more than 1000 ions per cm³ at 15 km away from the land surface [7].

3.2. Sunlight Including Ultraviolet

The photoelectric effect is the emission of electrons when a certain wavelength of light is shone onto a metallic surface. NAIs are generated by accepting these emitted electrons. The photoelectric effect may contribute less to the NAI generation as only some wavelength of lights shows the ability to emit electrons by lighting. One of the examples is the negative ion generator using an ultraviolet source to irradiate electrically conductive material, which was patented as early as 1964 (Patent No. US 3128378 A). In this patent, an ultraviolet lamp was used to irradiate metal materials, which photo-electrically eject electrons. The electrons then collide with air molecules and generate NAIs.

On the other hand, NAIs can be generated by a certain wavelength of lights through directly ionizing air molecules. For example, ultraviolet (UV) can be used to directly ionize air molecules to generate NAIs [5,8]. Actually, UV-mediated ionization is the dominant NAI sources in the above 60 km altitude of atmosphere [5]. These highly concentrated NAIs from UV in the upper layers of the atmosphere are diffused to the ground surface at low speeds. Ionization by UV radiation is not a major contributor of NAIs in the lower atmosphere due to the low dose of UV rays available in this layer [5]. Although reports showed that the UV rays significantly mediated the air ionization, little systematical study was carried out on the effect of artificial UV light on NAI generation. We carried out an experiment to investigate the contribution of UV light to the generation of NAIs (Figure 1; Table S1). The experiment was carried out in a growth chamber with 80 cm length × 80 cm width × 80 cm height and detailed description was provided in the Supplementary Experiment. NAIs were measured under UV light conditions with the normal light condition as a control. The data from three replicates of experiments showed that UV lights indeed promoted the NAI generation. In the chamber, the average NAI concentration was 344 ions/cm³ within one hour. The NAI concentration was increased to 825 ions/cm³ under UV light condition, significantly higher than the control (Figure 1A). Further observation showed that there were peaks of NAI generation within 8 min after UV lighting (Figure 1 B–D; Table S1). After the peaks, NAI concentration was kept at a relatively stable value but was still higher than the control. For all three replicates or each replicate, a nonparametric two-tailed Mann–Whitney U test was carried out as described in the Supplementary Experiment and the statistical analysis showed that NAI concentrations under UV lighting conditions in all three replicates were significantly higher than those under normal lighting conditions (CK) with $p < 0.00001$. The analysis further confirmed the promoting effect of UV lighting on NAI generation. Thus, our experiment showed that UV lighting could be used to generate NAIs. However, only low concentrations of NAIs were generated under our UV light conditions.

[Open in a separate window](#)

[Figure 1](#)

NAI generation by UV lighting. The experiment was carried out in a growth chamber with dimensions of 80 cm length, 80 cm width and 80 cm height. A 30 watts of UV light (UV-C, 100–280 nm) was provided by Safer Electric Ltd., Singapore. NAI concentration was measured by the DLY-4G-232 air ion counter (Kilter Electronic Institute Co., Ltd., Zhangzhou, Fujian Province, China). (A) The average NAI concentrations of control (CK, no UV light) and UV lighting. The NAI concentration under UV light was significantly higher than the control as indicated by Mann–Whitney U test at $p < 0.00001$. (B–D) Graphs to show the curves of NAI concentrations among three different replicates. The star “*” indicated that the NAI concentration under UV lighting was statistically higher than that in CK by Mann–Whitney U test at $p < 0.00001$. NAI: negative air ions.

3.3. Natural and Artificial Corona Discharge Including Thunderstorms and Lightning

The atmosphere surrounding the earth is subjected to a natural electric field and its intensity is continuously fluctuating under both local and global influences [9]. The local influences include geographical location and weather conditions such as thunderstorms, rain, fog, mist, and so on; the global facts refer to classical daily electric field variations [4]. When leaf points or branches of trees have a high potential difference from their surroundings in their electric fields, corona discharge (also called point discharge) occurs and NAIs may be released [5,10]. Generally, corona discharge occurs at the atmosphere conditions under high average electric fields [5]. For example, in a mountain area, high

electric fields and low atmospheric pressure promote the onset of corona discharge [11]. Thunderstorms and lightning will generate very high electric field conditions and corona discharge subsequently occurs. Therefore, NAIs will be released at a huge amount after thunderstorms and lightning. However, released NAIs will be gradually decayed with the discontinuous thunderstorms. In addition to thunderstorms and lightning, mist may also contribute to NAI generation. In a forest, electric field variations were observed during the mist formation and dissipation, which may trigger corona discharge and NAI generation [9].

Artificial corona discharge is an efficient way to generate NAIs. When a high negative voltage is applied to a conductor/electrode and generated electric field is high enough, corona discharge occurred [12,13]. If a charged conductor/electrode has a needle-type with a sharp point, the electric field around the tip will be significantly higher than other parts and air near the electrode can become ionized and NAIs are generated [14]. Intensity of corona discharge depends on the shape and size of the conductors as well as applied voltage. Irregular conductor, especially with a sharp point, gives rise to more corona than a smooth conductor and large-diameter conductors produce lower corona than small-diameter conductors; the higher the voltage applied, more NAIs are generated [14,15]. The closer the distance to corona point, the higher NAI concentration is detected as continuous generation of NAIs by corona discharge is related to a chain reaction process called an electron avalanche [16]. The application of artificial electric field and corona discharge on plants was carried out as early as the 1960s [17,18]. Bachman and Hademenos (1971) showed that under high voltage, artificially applied electrical fields near the pointed barley leaf tips were intensified [19] and as a result, corona discharge occurred and air ions and ozone were generated. Studies mainly focused on biological effects such as growth response, evaporation, and plant damage as well as the effects of generated ozone and NAIs on plant growth [11,18,19,20,21,22].

3.4. The Shearing Forces of Water (Lenard Effect)

The considerable numbers of NAIs are found under waterfalls or in the seashores. These NAIs are generated by Lenard effect. Lenard effect was also called spray electrification or waterfall effect and was first systematically studied by Philipp Lenard [23], who won the Nobel Prize for Physics in 1905 for his research on cathode rays and the discovery of many of their properties. The study showed that NAIs were generated from the surrounding air molecules by charging themselves negatively when water droplets collide with each other or with a wetted solid to form fine spray of drops. The study also showed that several factors may affect the degree of charge separation in spray processes and, therefore, may affect the generation and concentration of NAIs. These factors include water drop temperature, dissolved impurities, speed of the impinging air blast, and foreign impinging surfaces of droplets. Based on the “Lenard effect”, water shearing appliance has been designed to generate NAIs [24]. Water shearing produced only superoxide ions (O_2^-) which was bound to clusters of water molecules to form the structure $O_2^-(H_2O)_n$ [25], and was essentially regarded as a natural source of NAIs [24]. NAIs generated by the “Lenard effect” might improve erythrocyte deformability, thereby aerobic metabolism [24].

3.5. Plant-Based NAI Release under Normal Growth Conditions and by Pulse Electric Stimulation

Plants were reported to have the ability to generate NAIs under normal growth conditions and have been regarded as natural resources for NAI generation [26,27]. Different plants released different amounts of NAIs under natural growth conditions (Table 1). However, under normal growth conditions, plants released very low concentration of NAIs (<200 ions/cm³, Table 1). Bachman and Hademenos (1971) reported the NAI generation by applying high-voltage electric field to plants [19]. Later, Tikhonov et al. (2004) showed that plants could release huge amounts of NAIs under pulse electric field (PEF) stimulation [28]. Since then, several other studies were carried out to investigate

the effect of PEF stimulation on plant NAI generation ([Table 2](#)). Generally, under natural growth conditions, plants release less than 200 ions/cm³ ([Table 1](#)). However, after PEF stimulation, more than 3.5×10^6 ions/cm³ were detected ([Table 2](#)). Several parameters may affect the NAI release under PEF stimulation including plant species and output voltages in PEF stimulation ([Table 2](#)) as well as light intensity, temperature, pulse interval, and the pulse width of PEF [[28,29,30](#)]. Studies of plant morphology on NAI release showed that species with blade shapes generated higher concentration of NAIs [[27](#)]. All these studies may provide an alternative to artificially generate NAIs through a plant-based method.

Table 1

NAIs released by different plants under natural growth conditions without pulse electric field (PEF) stimulation.

[Open in a separate window](#)

* Specific distance may be related to plant size, please refer to the cited paper for more details. # Average values for all used plant species under 60 min of experiment.

Table 2

NAIs released by different plants under different treatments with PEF stimulation.

[Open in a separate window](#)

* Specific distance may be related to plant size, please refer to the cited paper for more details. kv: kilovoltage.

Besides the abovementioned 5 different sources of NAIs, some other ways also may generate NAIs. For example, the friction by rapidly moving great volumes of air over land may generate NAIs and storms have been also regarded as an energy source for NAI generation [[31](#)].

3.6. The Fate of Generated NAIs

As described above, we have reviewed the generation of various sources of NAIs and various factors affecting NAI generation. Here, we further explored the fate of generated NAIs. Generally, generated NAIs are not stable and will be gradually decayed. When NAIs combined with water molecules and form negative-ion clusters, their lifetime will be longer. For example, the half-life of negative oxygen ions $O_2^-(H_2O)_n$ by the Lenard effect is around 60 seconds, but the NAIs generated by corona discharge could survive only several seconds [[32](#)]. Such a trend in survival duration of these two different sources of NAIs was also observed by another study [[33](#)]. In the study, the lifetime of NAIs generated by the Lenard effect was several minutes, significantly longer than that of electrically generated NAIs. However, other studies showed that the lifetime of NAIs in clean air is around 100 s

([34] and references therein). In fact, several other factors including the concentrations of NAIs and aerosols, and the intensity of electric fields may affect lifetimes of NAIs [35]. This may explain why different lifetimes were reported by several different references as mentioned above.

4. Plants as a Source for Generating NAIs

Negative oxygen ions are the most commonly recognized NAIs. Reports showed that superoxide $O_2^{\bullet-}$ was a kind of NAIs [28,36,37,38]. Among NAIs generated by natural atmosphere and the Lenard effect (waterfall), superoxide $O_2^{\bullet-}$ are the major negative ions [37,38]. In plants, superoxide $O_2^{\bullet-}$ is mainly produced in the thylakoid membrane of photosystem I (PSI) [39]. Generation of superoxide $O_2^{\bullet-}$ within PSII has also been reported [40,41,42]. Some plant peroxidases are located in the apoplast by ionically or covalently bonding to cell wall polymers and these cell wall peroxidases might contribute to the generation of extracellular superoxide ions [43,44]. Different plant species might exhibit the difference in superoxide generation. For example, high rates of extracellular superoxide were generated in bryophytes and lichens [45]. In addition, generation of superoxide could be regulated by some external stimulation. For example, in chloroplasts, through treating plants with methyl viologen (MV) or paraquat under light conditions, it is possible to generate $O_2^{\bullet-}$ steadily by univalent reduction of MV or paraquat and subsequently transferring their electrons to oxygen [39,46,47]. Immediately induced and transient generation of extracellular superoxide anion was observed after adding salicylic acid (SA) to tobacco (*Nicotiana tabacum*) suspension culture [48].

Furthermore, the generation of superoxide in plant leaves can be improved by regulating gene expression. For instance, expression of the gene encoding Rho-related small G protein (ROP) 2 promoted superoxide generation in *Arabidopsis* leaf extracts [49]. Interestingly, the production of superoxide in plants could be improved by transgenic techniques. Rice *PsbSI* gene encodes a 22-kDa Photosystem (PS) II protein involving in nonphotochemical quenching (NPQ) of chlorophyll fluorescence. Disfunction of this gene by either transfer DNA (T-DNA) knockout or RNAi led to increased superoxide production [50]. The result provides a new way to improve plants in their superoxide release.

In addition to superoxide, other negative ions also contribute to the composition of plant NAIs. However, little is known on how these negative ions are generated and released in plants. Studies mainly focused on several factors that affect the release of NAIs in plants. Wang and Li (2009) investigated the effects of light intensities on NAI release in five plant species (*Aloe arborescens*, *Clivia miniata*, *Chlorophytum comosum*, *Opuntia brunnescens*, *Crassula portulacea*) [27]. Their data showed that the NAI release was significantly increased with the enhancement of light intensities for *Aloe arborescens*. However, for the remaining 4 plant species, less effect was observed and NAI release was not significantly affected by light intensities. Skromulis and Noviks (2012) surveyed the effects of meteorological factors on NAI release in a city and found that the highest level of NAI concentration was observed in morning sections and less air pollutants were detected in the duration [51]. Wu et al. (2017) analysed the NAI release of three plant species, *Crinum asiaticum*, *Narcissus tazetta*, and *Zephyranthes carinata*, under different light intensities and found that the stronger light might stimulate the plants to release more NAIs [30].

5. Composition of Negative Air Ions

Generally, NAIs are composed of multiple negatively charged molecules and these negative ions combine with several or up to 20 or 30 water molecules and form negative-ion clusters such as $CO_3^-(H_2O)_n$, $O^-(H_2O)_n$, and $O_3^-(H_2O)_n$. [24,52,53]. Mass spectrometric techniques were widely used to determine the composition of NAIs from various sources [54,55,56]. Early measurements suggested that the majority of lower tropospheric negative ions would be composed of O_2^- , CO_3^- , or NO_3^- and their $(H_2O)_n$ clusters as well as HSO_4^- core ions [5,57]. Studies showed that atmosphere

NAIs also included additional ions such as OH^- , NO_2^- , HCO_3^- and their water clusters [31,57].

[Figure 2](#) summarizes the composition from various sources of NAIs and the evolution of oxygen-based NAIs [31,37,52,53,56,57,58,59,60,61,62,63,64,65,66].

[Open in a separate window](#)

[Figure 2](#)

NAI species generated by different methods and evolution of oxygen-based NAIs. There were some differences in NAI compositions from different studies as their experimental conditions might vary. (A) NAI compositions. (B) The evolution of oxygen-based NAIs. The blue arrows indicate the NAI transformation processes. Please refer to the cited paper for more details [31,37,52,53,56,57,58,59,60,61,62,63,64,65,66].

Negative ion species generated by corona discharge have been identified through mass spectrometry by several experiments [55,66,67]. The majority of negative ions are CO_3^- and other negative ions include O^- , O_3^- , NO_3^- , and so on, which consist of less than 10% [52,67,68,69,70,71,72,73,74]. Reports from Nagato et al. (2006) showed that negative ion compositions are different at different reaction times of corona ionization as observed by mass spectrometry [74]. Based on their results, NO_3^- is the major ion followed by HCO_3^- and others. Both results from Shahin (1969) and Nagato et al. (2006) indicated that O_2^- is not dominant products of negative ions produced by corona discharge [67,74]. The majority of NAIs from corona discharge were listed in [Figure 2A](#).

Waterfalls-induced negative ions evolved from both O_2^- and O^- [56]. The OH^- concentration was significantly increased near waterfalls. These three ions, O_2^- , O^- , and OH^- , further evolved into other types of ions. As a result, the following 5 types of ions were regarded as the major negative air ions generated by waterfalls. They were listed in [Figure 2A](#). For the plant-based NAIs released by PEF stimulation, no reports were available on their ion composition.

Based on the studies mentioned above, NAIs may evolve from one NAI to another NAI. For example, the NAI O^- is formed when an oxygen molecule O_2 obtains an electron ([Figure 2B](#)). The NAI O^- may contribute to the formation of secondary NAIs by collision-aided electron attachment processes when other molecules exist in the same space [58]. As a result, other NAIs are generated such as O_2^- , CO_4^- , CO_3^- , OH^- , HCO_3^- , O_3^- , NO_3^- , and NO_2^- ([Figure 2B](#)). Parts et al. (2007) described a more complicated evolutionary process of NAIs [56]. In fact, the NAI evolution is related to the surrounding air composition. NAIs are continually changing as they collide with molecules in the air. Thus, NAIs are dynamic in their composition, which depends on ionization potential and electron affinity, the proton affinity, the dipole moment and the polarizability as well as the reactivity of the molecule [75].

6. Biological Effects of Negative Air Ions on Human/Animal Health and Microorganism Growth as well as Plant Development

Krueger and Reed (1976) reviewed key reports published in that period on the biological effects of NAIs [1]. They proposed the serotonin hypothesis of NAI biological actions. Serotonin is a very powerful and versatile neurohormone. It functions in inducing profound neurovascular, endocrinal, and metabolic effects in humans or animals. It plays important roles in basic patterns of life including sleep and mood regulation. Evidence has shown that NAIs could significantly reduce the level of serotonin in blood or brain, etc. [1,76,77]. Subsequent evidence showed that superoxide ions have been involved in the biological effects of NAIs [37] and in vitro experiments demonstrated that serotonin could be oxidized into tryptamine-4,5-dione by superoxide ions [78]. Thus, some of the biological effects of

NAIs were related to the reduction of serotonin. However, others reported no significant effect of NAIs on the concentration or turnover of serotonin in rats under their exposure conditions [35,79]. Bailey et al. (2018) carried out a comprehensive review on the effects of air ions on serotonin and other neurotransmitters and their analysis indicated some modest or strong evidence to support no effect hypothesis [35].

There are many references to report the possible biological effects [24,32,33,35,38,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100,101,102,103,104,105,106,107,108,109,110,111,112,113,114,115,116,117,118,119,120,121,122,123,124]. And some of them were listed in Table S2. The negative oxygen ion concentration exceeding 1000 ions/cm³ has been regarded as the threshold value for fresh air and the concentration should be higher for boosting the human immune system ([80] and references therein). Exposure to NAIs exhibit wide effects on animal/human health, anti-microorganisms and plant growth (Table S2). The effects of NAIs on human/animal health mainly focused on the cardiovascular and respiratory system as well as on mental health (Table S2). The effects of NAIs on the cardiovascular system included improving erythrocyte deformability and aerobic metabolism [24] as well as decreasing blood pressure [32,77,81,82]. However, no effect was also reported on blood pressure [83] or heart rate [84,85] and related data on cardiovascular function studies were not quantitatively evaluated [35]. On mental health, exposure to NAIs showed highly significant increase in performance of all tested tasks (mirror drawing, rotary pursuit, visual reaction time, and auditory) [86], alleviating symptoms of seasonal affective disorder (SAD) [87]. Similar effects of NAIs on relieving symptoms in mood disorders to antidepressant nonpharmacotherapy trials were observed [38]. NAIs also showed effective treatment of chronic depression [88]. On the contrary, no effect of NAIs on mental health was reported in other studies [89,90]. Comprehensive analysis on literatures showed no conclusive results on the potential therapeutic effects of NAIs on depression [35]. As for the effect of NAIs on respiratory function, exposure to negative or positive air ions does not appear to play an appreciable role in respiratory function [91]. In addition, some reports also showed the effects of NAIs on inhibiting cancer cells. For example, water-generated NAIs activated natural killer (NK) cell and inhibit carcinogenesis in mice [33]. The presence of NAIs is credited for increasing psychological health, productivity, and overall well-being [38,92,93]. Reports also showed that NAIs attached themselves to particles such as dust, mold spores, and other allergens [37]. As a result, people in the NAI atmosphere relieved symptoms of allergies to these particles. Generally, although some reports showed that air enriched with NAIs have multiple beneficial therapeutic effects in normalizing arterial pressure and blood rheology, supporting tissue oxygenation, easing stress conditions, and augmenting resistance to adverse factors [94], a systematic review suggested no consistent or reliable effects of NAIs on cardiovascular and respiratory system as well as on mental health [35].

Similarly, there are lots of research articles to present the effects of NAIs on the growth of microorganism (Table S2). The majority of the studies focused on bacteria and the presence of high concentration of NAIs inhibited the growth of bacteria. An early study showed that NAIs caused a significant amount of biological decay of the bacterium *Serratia marcescens* [95]. Exposure to NAIs showed inactivation or growth inhibition of the bacteria *E. coli*, *Candida albicans*, *Staphylococcus aureus*, *P. fluorescens* [96,97,98,99,100] and has a lethal effect on starved *Pseudomonas veronii* cells [101]. NAIs prevented 60% of tuberculosis (TB) infection and 51% of TB disease [102]. Except for the inhibition effect of NAIs on bacteria, reports also showed that NAIs inhibited the growth of fungi and viruses. For example, NAIs could inhibit the growth of *Penicillium notatum* [103]; the use of NAI generators reduced airborne transmission of Newcastle disease virus [104]. However, some controversial results have been reported on the growth inhibition of NAIs on microorganisms. For example, Fan et al. (2007) reported that NAIs had a very limited effect on *E. coli* that were inoculated on mung bean seed and apples [105]. In another report, seven bacterial species (*Staphylococcus aureus*,

Mycobacterium parafortuitum, *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, *Burkholderia cenocepacia*, *Bacillus subtilis*, and *Serratia marcescens*) were exposed to NAIs and only the growth *Mycobacterium parafortuitum* was inhibited [106].

Besides animals/humans and microorganisms, plants might also be affected by NAI exposure. After NAI treatment of *Avena sativa*, the fresh and dry weight were increased [107] and mean stem length and integral elongation were also increased [108]. Oxygen consumption was increased in barley seedlings after exposure to NAIs [109]. Plant height increased by 13–15% and dry weight increased by 18% under the growth environment with high concentrations of NAIs [110]. Lettuce plants exposed to NAIs showed vigorous growth with increased leaf area and fresh weight [111]. NAI treatment improved sprout growth and bacterial control during plant development [112]. NAIs have a positive effect on kale growth by improving fresh weight, macroelements and microelements [113]. However, no significant difference was observed between control and NAI-treated tobacco plants in nicotine and total alkaloid; the total nitrogen content in NAI-treated plants was slightly increased when compared with control plants [114].

7. Superoxide Involvement in the Biological Effects of Negative Air Ions

The oxygen molecule O_2 is a biradical with two unpaired electrons in different orbitals and is capable of accepting up to 4 electrons [125]. Superoxide $O_2^{\bullet-}$ is formed by accepting one electron. At least 4 oxidation states of oxygen have been identified including O_2 (dioxygen), O_2^+ (dioxygen cation), $O_2^{\bullet-}$ (superoxide) and O_2^{2-} (peroxide dianion) [126]. The monovalent reduction of O_2 gives birth to $O_2^{\bullet-}$, which serves as both a radicle and a negative ion; thus, the radicle sign “•” is added [116]. Therefore, superoxide $O_2^{\bullet-}$ is a kind of negative oxygen ions, which was documented by several reports [28,36,37,38,127]. Reports showed that superoxide ions are the major negative ions generated by natural atmosphere and the Lenard effect (waterfall) [37,38,128] and is more stable than other NAIs [25,127]. Superoxide $O_2^{\bullet-}$ was also detected in those NAIs that were generated by stimulating plant roots using PEF [28].

An early study showed the involvement of superoxide in the bacterial killing of *Staphylococcus albus* [117]. As commented by Rosenthal and Ben-Hur (1980) [36], the result was supported by the result that the bacterial killing of *Staphylococcus albus* by NAIs was protected by adding superoxide dismutase. However, reports showed that many other factors might also lead to the observed lethal effect of NAIs on bacteria [36]. Inhaled gaseous superoxide might improve the antinociceptive effect of analgesic agents and relieved the pathological signs of Parkinson’s disease [2,129]. Goldstein et al. (1992) also reported that the biological activity of NAIs may be dependent on superoxide [128], but it must be emphasized that the biological effects of NAIs are due to not only superoxide, but also other activated oxygen species [127,130,131]. On the other hand, it also needs to be mentioned that gas-phase superoxide ions are different from liquid-phase superoxide [37]; thus, the biological effects of the former might also be different from those of the latter.

8. The Release of Negative Air Ions as an Efficient Way to Remove Fine Particulates

PM is the major pollutant during haze episodes. Evidence showed that PM, especially $PM_{2.5}$, seriously affected human health. Based on the data from Lim et al. (2012), around 3.2 million people died in 2010 due to exposure to PM [132]. Annually, 2.1 million deaths occurred in the world due to the increasing levels of $PM_{2.5}$ [133,134]. PM mainly led to diseases related to both respiratory and cardiovascular systems. Fine PM such as $PM_{1.0}$ and $PM_{2.5}$ penetrate deeply into the lung, irritate and corrode the alveolar wall and, as a result, affect lung functions [135] and might also result in lung cancer [136]. However, the deposition of particles in the respiratory tract is only enhanced for particles

with many charges, which can only be achieved with an ionizer in an enclosed space. PM was also related to increased blood glucose, worse endothelial function, and incident cardiovascular disease events [137].

NAIs are unipolar ions and they will electrically charge PM, which will be deposited/precipitated more rapidly than uncharged ones as charged PM can attach to nearby surfaces, attach to each other and settle faster, or sink faster under the gravity [138,139,140,141,142] (available online: <https://www.epa.gov/indoor-air-quality-iaq/guide-air-cleaners-home>).

Tanaka et al. (1996) reported that NAIs reduced respirable and inhalable dust counts by 46% [143]. PM concentration was reduced by up to two orders of magnitude after 2 hours of treatment by NAI generator in an unoccupied office room (50 m³) [144]. Many other similar reports showed the high efficiency of NAIs in PM removal (Table 3; [138,140,145,146,147,148,149,150,151,152]). Evidence showed that many factors might affect PM reducing efficiency by NAIs (Table 3; [34,139,150,153]). For example, PM removal rate was related to particle concentrations, particle sizes, and the ventilation conditions and a model was raised to compute their relationship [139]. Studies also showed that wall surface significantly affected PM reducing efficiency [34]. In the chamber with the wood and polyvinyl chloride (PVC) wall surfaces, the NAI could remove PM particles substantially more effectively when compared with other wall materials such as wallpaper, stainless steel, and cement paint. One of the obvious disadvantages of ionizers is to emit ozone [154], which is a powerful oxidant and may seriously harm our health by long-term and/or high-dose exposure. Studies showed that many NAI generators including well-branded ionizers release ozone (Table 3; [141,145,155,156,157]). Another side effect is that the continuous emission of NAIs into an enclosed environment may lead to charge accumulation on insulating surfaces and then may cause electrostatic problems, especially when the humidity is low [138].

Table 3

The effect of NAIs on particulate matter (PM) removal.

[Open in a separate window](#)

9. Prospects and Future Works on Plant-Based NAI Generator

As mentioned above, plants could release a huge amount of negative air ions under PEF stimulation. This is an interesting topic and many more works should be carried out to figure out the mechanisms for plants to help generate NAIs. Evidence showed that light intensity significantly increased the plant-based NAI release by PEF stimulation [28]. During PEF stimulation, more leaf stomata opened wider which might contribute to a higher level of NAI release [28]. Generally, limited data are available on how PEF stimulation helps plants to generate very high concentrations of NAIs. No data were reported on what happens during PEF stimulation and how the stimulation may regulate the profile of gene expression. On the other hand, no reports were available on the composition of these NAIs released through stimulating plants by PEF treatment and their biological effects. Furthermore, only limited data were available on the application of these plant-based NAIs on reducing PM (available online: <https://patents.google.com/patent/CN203313745U/en?q=CN+203313745+U>). In a word, studies on the mechanism of plant-based release of NAIs by PEF stimulation are still under way. More efforts should be carried out on the comparison between these plant-based NAIs and other sources of NAIs and their difference in biological effects and PM reduction.

10. Conclusions

Some studies have suggested that NAIs had multiple health benefits on humans/animals, might inhibit the growth and/or kill some of microorganisms and promote plant development ([Figure 3](#)), but some of the results need to be further verified, some references might overestimate its benefits and no consistent or reliable evidence in therapeutic effects were achieved. However, to our knowledge, no data showed the harmful effects of NAIs on humans/animals. Superoxide ions are key members of NAIs and have been involved in the biological effects of NAIs by regulating the serotonin level and other biological actions but some reports showed no significant effect of NAIs on the concentration or turnover of serotonin. On the other hand, evidence showed that NAIs could high-efficiently remove PM including ultrafine PM ([Figure 3](#)), providing an alternative way to clean indoor air especially during haze episodes. We have also reviewed the plant-based NAI generation system by PEF stimulation. More efforts should be input into this system to further improve it and to use it as a high-efficient NAI generator and PM removal system.

[Open in a separate window](#)

[Figure 3](#)

A summary of the benefits of negative air ions on organisms and air cleaning.

Abbreviations

NAI	Negative air ion
PM	Particulate matter
PEF	Pulsed electric field
UV	Ultraviolet
SA	Salicylic acid
MV	Methyl viologen
KV	Kilovoltage
RANIs	Releasing amounts of negative ions
PVC	Polyvinyl chloride

Supplementary Materials

Supplementary materials can be found at <http://www.mdpi.com/1422-0067/19/10/2966/s1>.

[Click here for additional data file.](#) (792K, zip)

Author Contributions

S.R. supervised the study; S.-Y.J. conceived of the study, analyzed the data and wrote the article; A.M. carried out the experiments. S.R. supervised and complemented the writing. All authors read and approved the final manuscript.

Funding

This research was funded by the Singapore Millennium Foundation project “Bio-based strategies to improve indoor air quality by increasing negative air ion release”.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Krueger A.P., Reed E.J. Biological impact of small air ions. *Science*. 1976;193:1209–1213. doi: 10.1126/science.959834. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
2. Goldstein N., Arshavskaya T.V. Is atmospheric super oxide Vitally necessary? Accelerated death of animals in a quasi-neutral electric atmosphere. *Z. Naturforsch C*. 1997;52:396–404. doi: 10.1515/znc-1997-5-619. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
3. Laza V. The environment and gaseous ions. *Cent. Eur. J. Occup. Environ. Med*. 2000;6:3–10. [[Google Scholar](#)]
4. Hoppel W.A., Anderson R.V., Willet J.C. Atmospheric Electricity in the Planetary Boundary Layer, The Earth’s Electrical Environment. NAS Press; Washington, DC, USA: 1986. pp. 195–205. [[Google Scholar](#)]
5. Harrison R.G., Carslaw K.S. Ion-aerosol-cloud processes in the lower atmosphere. *Rev. Geophys*. 2003;41 doi: 10.1029/2002RG000114. [[CrossRef](#)] [[Google Scholar](#)]
6. Chalmers J.A. Atmospheric Electricity. 2nd ed. Pergamon Press; Oxford: UK; New York, NY, USA, 1967. [[Google Scholar](#)]
7. Ermakov V.I., Bazilevskaya G.A., Potrevsky P.E., Stozhkov Y.I. Ion balance equation in the atmosphere. *J. Geophys. Res*. 1997;102:23413–23419. doi: 10.1029/97JD01388. [[CrossRef](#)] [[Google Scholar](#)]
8. Yates A., Gray F.B., Misiaszek J.I., Wolman W. Air ions: Past problems and future directions. *Environ Int*. 1986;12:99–108. doi: 10.1016/0160-4120(86)90019-X. [[CrossRef](#)] [[Google Scholar](#)]
9. Borra J.P., Roos R.A., Renard D., Lazar H., Goldman A., Goldman M. Electrical and chemical consequences of point discharges in a forest during a mist and a thunderstorm. *J. Phys. D: Appl. Phys*. 1997;30:84–93. doi: 10.1088/0022-3727/30/1/011. [[CrossRef](#)] [[Google Scholar](#)]
10. Aubrecht L., Koller J., Stanek Z. Onset voltages of atmospheric corona discharges on plants. *Czech. J. Phys*. 2000;50:313–318. doi: 10.1007/BF03165903. [[CrossRef](#)] [[Google Scholar](#)]
11. Aubrecht L., Stanek Z., Koller J. Corona discharge on coniferous trees—Spruce and pine. *Europhys. Lett*. 2001;53:304–390. doi: 10.1209/epl/i2001-00153-2. [[CrossRef](#)] [[Google Scholar](#)]
12. Altamimi G., Illias H.A., Mokhtar N., Mokhlis H., Bakar A.H.A. Corona discharges under various types of electrodes; Proceedings of the 2014 IEEE International Conference on Power and Energy (PECon); Kuching Sarawak, Malaysia. 1–3 December 2014; pp. 5–8. [[CrossRef](#)] [[Google Scholar](#)]
13. Ogar V.N., Bendor S.A., James A.E. Analysis of corona effect on transmission line. *Am. J. Eng. Res*. 2017;6:75–87. [[Google Scholar](#)]
14. Berendt A., Budnarowska M., Mizeraczyk J. DC negative corona discharge characteristics in air flowing transversely and longitudinally through a needle-plate electrode gap. *J. Electrostat*. 2018;92:24–30. doi: 10.1016/j.elstat.2018.01.006. [[CrossRef](#)] [[Google Scholar](#)]
15. Sharma S., Goel K., Gupta A., Kumar H. Corona effects on eHv AC transmission lines. *Int. J. Sci. Res. Eng. Technol*. 2012;1:160–164. [[Google Scholar](#)]

16. Manabe Y., Shimazaki T. Formation mechanism of surface corona on dielectric plates under negative impulse voltage in atmospheric air. *IEEE Trans. Dielectr. Electr. Insul.* 2004;11:631–637. doi: 10.1109/TDEI.2004.1324353. [[CrossRef](#)] [[Google Scholar](#)]
17. Murr L.E. Plant growth response in a simulated electric field-environment. *Nature.* 1963;200:490–491. doi: 10.1038/200490b0. [[CrossRef](#)] [[Google Scholar](#)]
18. Murr L.E. Biophysics of plant growth in an electrostatic field. *Nature.* 1965;206:467–470. doi: 10.1038/206467a0. [[CrossRef](#)] [[Google Scholar](#)]
19. Bachman C.H., Hademenos D.G. Ozone and air ions accompanying biological applications of electric fields. *J. Atmos. Terr. Phys.* 1971;33:497–505. doi: 10.1016/0021-9169(71)90153-X. [[CrossRef](#)] [[Google Scholar](#)]
20. Krueger A.P., Strubbe A.E., Yost M.G., Reed E.J. Electric fields, small air ions and biological effects. *Int. J. Biometeorol.* 1978;22:202–212. doi: 10.1007/BF01555399. [[CrossRef](#)] [[Google Scholar](#)]
21. Hart F.X., Schottenfeld R.S. Evaporation and plant damage in electric fields. *Int. J. Biometeor.* 1979;23:63–68. doi: 10.1007/BF01553377. [[CrossRef](#)] [[Google Scholar](#)]
22. Hart F.X., Mudano M.J., Atchley A.A. Plant damage produced by the passage of low level direct current. II. Further observations and possible mechanisms. *Int. J. Biometeor.* 1981;25:151–159. doi: 10.1007/BF02184463. [[CrossRef](#)] [[Google Scholar](#)]
23. Lenard P. Über die Electrizedität der Wasserfälle. *Ann. Phys.* 1892;46:584–636. doi: 10.1002/andp.18922820805. [[CrossRef](#)] [[Google Scholar](#)]
24. Iwama H. Negative air ions created by water shearing improve erythrocyte deformability and aerobic metabolism. *Indoor Air.* 2004;14:293–297. doi: 10.1111/j.1600-0668.2004.00254.x. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
25. Kosenko E.A., Kaminsky Y.G., Stavrovskaya I.G., Sirota T.V., Kondrashova M.N. The stimulatory effect of negative air ions and hydrogen peroxide on the activity of superoxide dismutase. *FEBS Lett.* 1997;410:309–312. doi: 10.1016/S0014-5793(97)00651-0. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
26. Nemeryuk G.E. Migration of salts into the atmosphere during transpiration. *Sov. Plant Physiol.* (Engl. Transl.) 1970;17:560–566. [[Google Scholar](#)]
27. Wang J., Li S.H. Changes in negative air ions concentration under different light intensities and development of a model to relate light intensity to directional change. *J. Environ. Manag.* 2009;90:2746–2754. doi: 10.1016/j.jenvman.2009.03.003. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
28. Tikhonov V.P., Tsvetkov V.D., Litvinova E.G., Sirota T.V., Kondrashova M.N. Generation of negative air ions by plants upon pulsed stimulation applied to soil. *J. Plant Physiol.* 2004;51:414–419. doi: 10.1023/B:RUPP.0000028690.74805.e2. [[CrossRef](#)] [[Google Scholar](#)]
29. Zhu M., Zhang J., You Q., Banuelos G.S., Yu Z., Li M., Bazai N.A., Tahir M. Bio-generation of negative air ions by grass upon electrical stimulation applied to lawn. *Fresen. Environ. Bull.* 2016;25:2071–2078. [[Google Scholar](#)]
30. Wu R., Zheng J., Sun Y., Wang Q., Deng C., Ye D. Research on generation of negative air ions by plants and stomatal characteristics under pulsed electrical field stimulation. *Int. J. Agric. Biol.* 2017;19:1235–1245. doi: 10.17957/IJAB/15.0431. [[CrossRef](#)] [[Google Scholar](#)]
31. Guerra L. A new instrument for the measurement of air ions. *Int. J. Biometeorol.* 1988;32:11–16. doi: 10.1007/BF01623986. [[CrossRef](#)] [[Google Scholar](#)]

32. Iwama H., Ohmizo H., Furuta S., Ohmori S., Watanabe K., Kaneko T., Tsutsumi K. Inspired superoxide anions attenuate blood lactate concentrations in postoperative patients. *Crit. Care Med.* 2002;30:1246–1249. doi: 10.1097/00003246-200206000-00014. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
33. Yamada R., Yanoma S., Akaike M., Tsuburaya A., Sugimasa Y., Takemiya S., Motohashi H., Rino Y., Takanashi Y., Imada T. Water-generated negative air ions activate NK cell and inhibit carcinogenesis in mice. *Cancer Lett.* 2006;239:190–197. doi: 10.1016/j.canlet.2005.08.002. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
34. Wu C.C., Lee G.W.M., Cheng P., Yang S., Yu K.P. Effect of wall surface materials on deposition of particles with the aid of negative air ions. *Aerosol Sci.* 2006;37:616–630. doi: 10.1016/j.jaerosci.2005.05.018. [[CrossRef](#)] [[Google Scholar](#)]
35. Bailey W.H., Williams A.L., Leonhard M.J. Exposure of laboratory animals to small air ions: A systematic review of biological and behavioral studies. *BioMed. Eng. online.* 2018;17:72. doi: 10.1186/s12938-018-0499-z. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
36. Rosenthal I., Ben-Hur E. Superoxide involvement in negative air ion effects. *Nature.* 1980;288:739–740. doi: 10.1038/288739a0. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
37. Goldstein N. Reactive oxygen species as essential components of ambient air. *Biochemistry.* 2002;67:161–170. doi: 10.1023/A:1014405828966. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
38. Pino O., Ragione F.L. There's something in the air: Empirical evidence for the effects of negative air ions (NAI) on psychophysiological state and performance. *Res. Psychol. Behav Sci.* 2013;1:48–53. [[Google Scholar](#)]
39. Scarpeci T.E., Zanol M.I., Carrillo N., Mueller-Roeber B., Valle E.M. Generation of superoxide anion in chloroplasts of *Arabidopsis thaliana* during active photosynthesis: A focus on rapidly induced genes. *Plant Mol. Biol.* 2008;66:361–378. doi: 10.1007/s11103-007-9274-4. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
40. Ananyev G.M., Renger G., Wacker U., Klimov V.V. Photoreduction of superoxide radicals and the superoxide dismutase activity of Photosystem II. The possible involvement of cytochrome b559. *Photosynth. Res.* 1994;41:327–338. doi: 10.1007/BF00019410. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
41. Cleland R.E., Grace S.C. Voltammetric detection of superoxide production by photosystem II. *FEBS Lett.* 1999;457:348–352. doi: 10.1016/S0014-5793(99)01067-4. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
42. Pospíšil P., Šnyrychova I., Kruk J., Strzałka K., Nauš J. Evidence that cytochrome b559 is involved in superoxide production in photosystem II: Effect of synthetic short-chain plastoquinone in a cytochrome b559 tobacco mutant. *Biochem. J.* 2006;397:321–327. doi: 10.1042/BJ20060068. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
43. Allan A., Fluhr R. Two distinct sources of elicited reactive oxygen species in tobacco epidermal cells. *Plant Cell.* 1997;9:1559–1577. doi: 10.1105/tpc.9.9.1559. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
44. Bolwell G.P., Davies D.R., Gerrish C., Auh C.-K., Murphy T.M. Comparative biochemistry of the oxidative burst produced by rose and french bean cells reveals two distinct mechanisms. *Plant Physiol.* 1998;116:1379–1385. doi: 10.1104/pp.116.4.1379. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

45. Minibayeva F., Beckett R.P. High rates of extracellular superoxide production in bryophytes and lichens, and an oxidative burst in response to rehydration following desiccation. *New Phytol.* 2001;152:333–343. doi: 10.1046/j.0028-646X.2001.00256.x. [[CrossRef](#)] [[Google Scholar](#)]
46. Asada K., Takahashi M. Production and Scavenging of Active Oxygens in Photosynthesis. In: Kyle D.J., Osmond C.B., Arntzen D.J., editors. *Photoinhibition*. Elsevier/North Holland Biomedical Press; Amsterdam, The Netherlands: 1987. pp. 227–280. [[Google Scholar](#)]
47. Halliwell B., Gutteridge J.M.C. *Free Radicals in Biology and Medicine*. Oxford University Press; Oxford, UK: 1989. [[Google Scholar](#)]
48. Kawano T., Sahashi N., Takahashi K., Uozumi N., Muto S. Salicylic acid induces extracellular superoxide generation followed by an increase in cytosolic calcium ion in tobacco suspension culture: The earliest events in salicylic acid signal transduction. *Plant Cell Physiol.* 1998;39:721–730. doi: 10.1093/oxfordjournals.pcp.a029426. [[CrossRef](#)] [[Google Scholar](#)]
49. Park J., Gu Y., Lee Y., Yang Z., Lee Y. Phosphatidic acid induces leaf cell death in Arabidopsis by activating the Rho-related small G protein GTPase-mediated pathway of reactive oxygen species generation. *Plant Physiol.* 2004;134:129–136. doi: 10.1104/pp.103.031393. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
50. Zulfugarov I.S., Tovuu A., Eu Y.J., Dogsom B., Poudyal R.S., Nath K., Hall M., Banerjee M., Yoon U.C., Moon Y.H., et al. Production of superoxide from Photosystem II in a rice (*Oryza sativa* L.) mutant lacking PsbS. *BMC Plant Biol.* 2014;14:242. doi: 10.1186/s12870-014-0242-2. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
51. Skromulis A., Noviks G. Atmospheric light air ion concentrations and related meteorologic factors in Rezekne city, Latvia. *J. Environ. Biol.* 2012;33:455–462. [[PubMed](#)] [[Google Scholar](#)]
52. Gravendeel B., de Hoog F.J. Clustered negative ions in atmospheric negative corona discharges in the Trichel regime. *J. Phys. B: Atom. Mol. Phys.* 1999;20:6337–6361. doi: 10.1088/0022-3700/20/23/025. [[CrossRef](#)] [[Google Scholar](#)]
53. Luts A., Parts T. Evolution of negative small air ions at two different temperatures. *J. Atmos. Sol-Terr. Phy.* 2002;64:763–774. doi: 10.1016/S1364-6826(02)00076-7. [[CrossRef](#)] [[Google Scholar](#)]
54. Dillard J.G. Negative ion mass spectrometry. *Chem. Rev.* 1973;73:589–643. doi: 10.1021/cr60286a002. [[CrossRef](#)] [[Google Scholar](#)]
55. Skalny J.D., Orszagh J., Mason N.J., John R., Yolanda A.G., Whitmore T.D. Mass spectrometric study of negative ions extracted from point to plane negative corona discharge in ambient air at atmospheric pressure. *Int. J. Mass Spectrom.* 2008;272:12–21. doi: 10.1016/j.ijms.2007.12.012. [[CrossRef](#)] [[Google Scholar](#)]
56. Parts T.E., Luts A., Laakso L., Hirsikko A., Gronholm T., Kulmala M. Chemical composition of waterfall-induced air ions: Spectrometry vs. simulations. *Boreal Environ. Res.* 2007;12:409–420. [[Google Scholar](#)]
57. Cabane M., Playe P. Mass spectra of negative ions in air-like gas mixtures at atmospheric pressure. *J. Aerosol Sci.* 1980;11:475–482. doi: 10.1016/0021-8502(80)90120-2. [[CrossRef](#)] [[Google Scholar](#)]
58. Lin H.F., Lin J.M. Generation and determination of negative air ions. *J. Anal. Test.* 2017;1:6. doi: 10.1007/s41664-017-0007-7. [[CrossRef](#)] [[Google Scholar](#)]
59. Goldman M., Goldman A., Sigmond R.S. The corona discharge, its properties and specific uses. *Pure Appl. Chem.* 1985;57:1353–1362. doi: 10.1351/pac198557091353. [[CrossRef](#)] [[Google Scholar](#)]

60. Sauers I., Harman G. A mass spectrometric study of positive and negative ion formation in an sF6 corona. Part II: Influence of water and sF6 neutral by-products. *J. Phys. D: Appl. Phys.* 1992;25:774–782. doi: 10.1088/0022-3727/25/5/006. [[CrossRef](#)] [[Google Scholar](#)]
61. Stano M., Safonov E., Kucera M., Matejcik S. Ion mobility spectrometry study of negative corona discharge in oxygen/nitrogen mixtures. *Chemicke Listy.* 2008;102:s1414–s1417. [[Google Scholar](#)]
62. Manninen H.E., Franchin A., Schobesberger S., Hirsikko A., Hakala J., Skromulis A., Kangasluoma J., Ehn M., Junninen H., Mirme A., et al. Characterization of corona-generated ions used in a Neutral cluster and Air Ion Spectrometer (NAIS) Atmos. Meas. Tech. 2011;4:2767–2776. doi: 10.5194/amt-4-2767-2011. [[CrossRef](#)] [[Google Scholar](#)]
63. Sekimoto K., Takayama M. Collision-induced dissociation analysis of negative atmospheric ion adducts in atmospheric pressure corona discharge ionization mass spectrometry. *J. Am. Soc. Mass Spectrom.* 2013;24:780–788. doi: 10.1007/s13361-013-0576-2. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
64. Laakso L., Hirsikko A., Grönholm T., Kulmala M., Luts A., Parts T.E. Waterfalls as sources of small charged aerosol particles. *Atmos. Chem. Phys.* 2006;6:9297–9314. doi: 10.5194/acpd-6-9297-2006. [[CrossRef](#)] [[Google Scholar](#)]
65. Kolarz P., Gaisberger M., Madl P., Hofmann W., Ritter M., Hartl A. Characterization of ions at alpine waterfalls. *atmos. Chem. Phys.* 2012;12:3687–3697. doi: 10.5194/acp-12-3687-2012. [[CrossRef](#)] [[Google Scholar](#)]
66. Skalny J.D., Horvath G., Mason N.L. Mass spectrometric analysis of small negative ions ($e/m < 100$) produced by trichel pulse negative corona discharge fed by ozonised air. *J. Optoelectron. Adv. Mater.* 2007;9:887–893. [[Google Scholar](#)]
67. Shahin M.M. Nature of charge carriers in negative coronas. *Appl. Opt.* 1969;8:106–110. doi: 10.1364/AO.8.S1.000106. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
68. Gardiner P.S., Moruzzi J.L., Craggs J.D. Negative ions in Trichel corona in carbon dioxide and oxygen. *J. Phys. D: Appl. Phys.* 1978;11:237–244. doi: 10.1088/0022-3727/11/3/009. [[CrossRef](#)] [[Google Scholar](#)]
69. Gardiner P.S., Craggs J.D. Negative ions in Trichel corona in air. *J. Phys. D: Appl. Phys.* 1977;10:1003–1009. doi: 10.1088/0022-3727/10/7/006. [[CrossRef](#)] [[Google Scholar](#)]
70. Sakata S., Okada T. Effect of humidity on hydrated cluster-ion formation in a clean room corona discharge neutralizer. *J. Aerosol. Sci.* 1994;25:879–885. doi: 10.1016/0021-8502(94)90054-X. [[CrossRef](#)] [[Google Scholar](#)]
71. Skalnya J.D., Mikoviny T., Matejcik S., Mason N.J. An analysis of mass spectrometric study of negative ions extracted from negative corona discharge in air. *Int. J. Mass Spectrom.* 2004;233:317–324. doi: 10.1016/j.ijms.2004.01.012. [[CrossRef](#)] [[Google Scholar](#)]
72. Hill C.A., Thomas C.L. A pulsed corona discharge switchable high resolution ion mobility spectrometer-mass spectrometer. *Analyst.* 2003;128:55–60. doi: 10.1039/b207558j. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
73. Ross S.K., Bell A.J. Reverse flow continuous corona discharge ionisation applied to ion mobility spectrometry. *Int. J. Mass Spectrom.* 2002;218:L1–L6. doi: 10.1016/S1387-3806(02)00693-0. [[CrossRef](#)] [[Google Scholar](#)]
74. Nagato K., Matsui Y., Miyata T., Yamauchi T. An analysis of the evolution of negative ions produced by a corona ionizer in air. *Int. J. Mass Spectrom.* 2006;248:142–147. doi: 10.1016/j.ijms.2005.12.001. [[CrossRef](#)] [[Google Scholar](#)]

75. Charry J.M., Kavet R. *Air Ions: Physical and Biological Aspects*. CRC Press; Boca Raton, FL, USA: 1987. [[Google Scholar](#)]
76. Diamond M.C., Connor J.R., Jr., Orenberg E.K., Bissell M., Yost M., Krueger A. Environmental influences on serotonin and cyclic nucleotides in rat cerebral cortex. *Science*. 1980;210:652–654. doi: 10.1126/science.6254145. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
77. Ryushi T., Kita I., Sakurai T., Yasumatsu M., Isokawa M., Aihara Y., Hama K. The effect of exposure to negative air ions on the recovery of physiological responses after moderate endurance exercise. *Int. J. Biometeorol.* 1998;41:132–136. doi: 10.1007/s004840050066. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
78. Wrona M.Z., Dryhurst G. Oxidation of serotonin by superoxide radical: Implications to neurodegenerative brain disorders. *Chem. Res. Toxicol.* 1998;11:639–650. doi: 10.1021/tx970185w. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
79. Bailey W.H., Charry J.M. Acute exposure of rats to air ions: Effects on the regional concentration and utilization of serotonin in brain. *Bioelectromagnetics*. 1987;8:173–181. doi: 10.1002/bem.2250080208. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
80. Lazzerini F.T., Orlando M.T., De Prá W. Progress of negative air ions in health tourism environments applications. *Bol. Soc. Esp. Hidrol. Méd.* 2018;33:27–46. doi: 10.23853/bsehm.2018.0450. [[CrossRef](#)] [[Google Scholar](#)]
81. Ju K., Kubo T. Power spectral analysis of autonomic nervous activity in spontaneously hypertensive rats. *Biomed. Sci. Instrum.* 1997;33:338–343. [[PubMed](#)] [[Google Scholar](#)]
82. Wiszniewski A., Suchanowski A., Wielgomas B. Effects of Air-Ions on human circulatory indicators. *Pol. J. Environ. Stud.* 2014;23:521–531. [[Google Scholar](#)]
83. Watanabe I., Noro H., Ohtsuka Y., Mano Y., Agishi Y. Physical effects of negative air ions in a wet sauna. *Int. J. Biometeorol.* 1997;40:107–112. doi: 10.1007/s004840050028. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
84. Bachman C.H., McDonald R.D., Lorenz P.J. Some physiological effects of measured air ions. *Int. J. Biometeorol.* 1965;9:127–139. doi: 10.1007/BF02188468. [[CrossRef](#)] [[Google Scholar](#)]
85. McDonald R.D., Bachman C.H., Lorenz P.J. Some physiological effects of air ion treatment without ion inhalation. *Int. J. Biometeorol.* 1965;9:141–147. doi: 10.1007/BF02188469. [[CrossRef](#)] [[Google Scholar](#)]
86. Hawkins L.H., Barker T. Air ions and human performance. *Ergonomics*. 1978;21:273–278. doi: 10.1080/00140137808931724. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
87. Bowers B., Flory R., Ametepe J., Staley L., Patrick A., Carrington H. Controlled trial evaluation of exposure duration to negative air ions for the treatment of seasonal affective disorder. *Psychiatry Res.* 2018;259:7–14. doi: 10.1016/j.psychres.2017.08.040. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
88. Goel N., Terman M., Terman J.S., Macchi M.M., Stewart J.W. Controlled trial of bright light and negative air ions for chronic depression. *Psychol. Med.* 2005;35:945–955. doi: 10.1017/S0033291705005027. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
89. Hedge A., Collis M.D. Do negative air ions affect human mood and performance? *Ann. Occup. Hyg.* 1987;31:285–290. [[PubMed](#)] [[Google Scholar](#)]
90. Reilly T., Stevenson I.C. An investigation of the effects of negative air ions on responses to submaximal exercise at different times of day. *J. Hum. Ergol.* 1993;22:1–9. [[PubMed](#)] [[Google Scholar](#)]

91. Alexander D.D., Bailey W.H., Perez V., Mitchell M.E., Su S. Air ions and respiratory function outcomes: A comprehensive review. *J. Negat. Results Biomed.* 2013;12:14. doi: 10.1186/1477-5751-12-14. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
92. Ogungbe A.S., Akintoye O.H., Idowu B.A. Effects of gaseous ions on the environment and human performance. *Trends Appl. Sci. Res.* 2011;6:130–133. doi: 10.3923/tasr.2011.130.133. [[CrossRef](#)] [[Google Scholar](#)]
93. Perez V., Alexander D.D., Bailey W.H. Air ions and mood outcomes: A review and meta-analysis. *BMC Psychiatry.* 2013;13:29. doi: 10.1186/1471-244X-13-29. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
94. Voeikov V.L. Reactive oxygen species--(ROS) pathogens or sources of vital energy? Part 1. ROS in normal and pathologic physiology of living systems. *J. Altern. Complement. Med.* 2006;12:111–118. doi: 10.1089/acm.2006.12.111. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
95. Phillips G., Harris G.J., Jones M.W. Effects of air ions on bacterial aerosols. *Int. J. Biometeorol.* 1964;8:27–37. doi: 10.1007/BF02186925. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
96. Tyagi A.K., Nirala B.K., Malik A., Singh K. The effect of negative air ion exposure on *Escherichia coli* and *Pseudomonas fluorescens*. *J. Environ. Sci. Health A Tox. Hazard Subst. Environ. Eng.* 2008;43:694–699. doi: 10.1080/10934520801959831. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
97. Shargawi J.M., Theaker E.D., Drucker D.B., MacFarlane T., Duxbury A.J. Sensitivity of *Candida albicans* to negative air ion streams. *J. Appl. Microbiol.* 1999;87:889–897. doi: 10.1046/j.1365-2672.1999.00944.x. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
98. Dobrynin D., Friedman G., Fridman A., Starikovskiy A. Inactivation of bacteria using DC corona discharge: Role of ions and humidity. *New J. Phys.* 2011:103033. doi: 10.1088/1367-2630/13/10/103033. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
99. Noyce J.O., Hughes J.F. Bactericidal effects of negative and positive ions generated in nitrogen on *Escherichia coli*. *J. Electrostatics.* 2002;54:179–187. doi: 10.1016/S0304-3886(01)00179-6. [[CrossRef](#)] [[Google Scholar](#)]
100. Timoshkin I.V., Maclean M., Wilson M.P., Given M.J., MacGregor S.J., Wang T., Anderson J.G. Bactericidal effect of corona discharges in atmospheric air. *IEEE Trans. Plasma Sci.* 2012;40:2322–2333. doi: 10.1109/TPS.2012.2193621. [[CrossRef](#)] [[Google Scholar](#)]
101. Noyce J.O., Hughes J.F. Bactericidal effects of negative and positive ions generated in nitrogen on starved *Pseudomonas veronii*. *J. Electrostatics.* 2003;57:49–58. doi: 10.1016/S0304-3886(02)00110-9. [[CrossRef](#)] [[Google Scholar](#)]
102. Escombe A.R., Moore D.A., Gilman R.H., Navincopa M., Ticona E., Mitchell B., Noakes C., Martínez C., Sheen P., Ramirez R., et al. Upper-room ultraviolet light and negative air ionization to prevent tuberculosis transmission. *PLoS Med.* 2009;6:e43. doi: 10.1371/journal.pmed.1000043. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
103. Pratt R., Barnard R.W. Some effects of ionized air on *Penicillium notatum*. *J. Am. Pharm. Assoc. Sci.* 1960;49:643–646. doi: 10.1002/jps.3030491004. [[CrossRef](#)] [[Google Scholar](#)]
104. Mitchell B.W., King D.J. Effect of negative air ionization on airborne transmission of newcastle disease virus. *Avian. Dis.* 1994;38:725–732. doi: 10.2307/1592107. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

105. Fan X., Fett W.F., Mitchell B.W. Effect of negative air ions on *Escherichia coli* ATCC 25922 inoculated onto mung bean seed and apple fruit. *J. Food Prot.* 2007;70:204–208. doi: 10.4315/0362-028X-70.1.204. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
106. Fletcher L.A., Gaunt L.F., Beggs C.B., Shepherd S.J., Sleigh P.A., Noakes C.J., Kerr K.G. Bactericidal action of positive and negative ions in air. *BMC Microbiol.* 2007;7:32. doi: 10.1186/1471-2180-7-32. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
107. Krueger A.P., Kotaka S., Andriese P.C. Some observations on the physiological effects of gaseous ions. *Int. J. Biometeorol.* 1962;6:33–48. doi: 10.1007/BF02187011. [[CrossRef](#)] [[Google Scholar](#)]
108. Krueger A.P., Kotaka S., Andriese P.C. Studies on the Effects of Gaseous Ions on Plant Growth I. The influence of positive and negative air ions on the growth of *Avena sativa*. *J. Gen Physiol.* 1962;45:879–895. doi: 10.1085/jgp.45.5.879. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
109. Kotaka S., Krueger A.P., Andriese P.C., Nishizawa K., Ohuchi T., Takenobu M., Kozure Y. Air ion effects on the oxygen consumption of barley seedlings. *Nature.* 1965;208:1112–1113. doi: 10.1038/2081112a0. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
110. Elkiey T.M., Bhartendu S., Barthakur N. Air ion effect on respiration and photosynthesis of barley and *Antirrhinum majus*. *Int. J. Biometeorol.* 1985;29:285–292. doi: 10.1007/BF02189659. [[CrossRef](#)] [[Google Scholar](#)]
111. Song M.J., Kang T.H., Han C.S., Oh M.M. Air anions enhance lettuce growth in plant factories. *Hortic. Environ. Biotechnol.* 2014;55:293–298. doi: 10.1007/s13580-014-1016-3. [[CrossRef](#)] [[Google Scholar](#)]
112. Song M.J., Jeon Y.M., Oh M.M. Growth of and bacterial counts on several edible sprouts exposed to spray ionization. *Protected Hort. Plant Fac.* 2015;24:45–50. doi: 10.12791/KSBEC.2015.24.1.045. [[CrossRef](#)] [[Google Scholar](#)]
113. Lee S.R., Kang T.H., Han C.S., Oh M.M. Air anions improve growth and mineral content of kale in plant factories. *Hortic. Environ. Biotechnol.* 2015;56:462–471. doi: 10.1007/s13580-015-0035-z. [[CrossRef](#)] [[Google Scholar](#)]
114. Barthakur N.N., Arnold N.P. Growth and certain chemical constituents of tobacco plants exposed to air ions. *Int. J. Biometeorol.* 1988;32:78–80. doi: 10.1007/BF01044897. [[CrossRef](#)] [[Google Scholar](#)]
115. Sirota T.V., Safronova V.G., Amelina A.G., Mal'tseva V.N., Avkhacheeva N.V., Sofin A.D., Ianin V.A., Mubarakshina E.K., Romanova L.K., Novoselov VI. Effect of negative air ions on respiratory organs and blood. *Biofizika.* 2008;53:886–893. doi: 10.1134/S0006350908050242. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
116. Wallner P., Kundi M., Panny M., Tappler P., Hutter H.P. Exposure to air ions in indoor environments: Experimental study with healthy adults. *Int. J. Environ. Res. Public Health.* 2015;12:14301–14311. doi: 10.3390/ijerph121114301. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
117. Kellogg E.W., Yost M.G., Barthakur N., Kreuger A.P. Superoxide involvement in the bactericidal effects of negative air ions on *Staphylococcus albus*. *Nature.* 1979;281:400–401. doi: 10.1038/281400a0. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
118. Gabbay J., Bergerson O., Levi N., Brenner S., Eli I. Effect of ionization on microbial air pollution in the dental clinic. *Environ. Res.* 1990;52:99–106. doi: 10.1016/S0013-9351(05)80154-9. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

119. Seo K.H., Mitchell B.W., Holt P.S., Gast R.K. Bactericidal effects of negative air ions on airborne and surface *Salmonella enteritidis* from an artificially generated aerosol. *J. Food Prot.* 2001;64:113–116. doi: 10.4315/0362-028X-64.1.113. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
120. Fan L., Song J., Hildebrand P.D., Forney C.F. Interaction of ozone and negative air ions to control micro-organisms. *J. Appl. Microbiol.* 2002;93:144–148. doi: 10.1046/j.1365-2672.2002.01683.x. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
121. Kerr K.G., Beggs C.B., Dean S.G., Thornton J., Donnelly J.K., Todd N.J., Sleigh P.A., Qureshi A., Taylor C.C. Air ionisation and colonization / infection with methicillin-resistant *Staphylococcus aureus* and *Acinetobacter* species in an intensive care unit. *Intensive Care Med.* 2006;32:315–317. doi: 10.1007/s00134-005-0002-8. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
122. Kampmann Y., Klingshirn A., Kloft K., Kreyenschmidt J. The application of ionizers in domestic refrigerators for reduction in airborne and surface bacteria. *J. Appl. Microbiol.* 2009;107:1789–1798. doi: 10.1111/j.1365-2672.2009.04359.x. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
123. Tyagi A.K., Malik A. Antimicrobial action of essential oil vapours and negative air ions against *Pseudomonas fluorescens*. *Int. J. Food Microbiol.* 2010;143:205–210. doi: 10.1016/j.ijfoodmicro.2010.08.023. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
124. Zhou P., Yang Y., Huang G., Lai A.C.K. Numerical and experimental study on airborne disinfection by negative ions in air duct flow. *Build. Environ.* 2018;127:204–210. doi: 10.1016/j.buildenv.2017.11.006. [[CrossRef](#)] [[Google Scholar](#)]
125. Alisa P. Oxygen Toxicity and Free Radical Injury. In: Marks A.D., Liebermann M.A., editors. *Mark's Basic Medical Biochemistry: A Clinical Approach*. 4th ed. Lippincott Williams & Wilkins; Philadelphia, PA, USA: 2013. pp. 437–456. [[Google Scholar](#)]
126. Hayyan M., Hashim M.A., AlNashef I.M. Superoxide ion: Generation and chemical implications. *Chem Rev.* 2016;116:3029–3085. doi: 10.1021/acs.chemrev.5b00407. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
127. Goldstein N.I., Goldstein R.N., Merzlyak M.N. Negative air ions as a source of superoxide. *Int. J. Biometeorol.* 1992;36:118–122. doi: 10.1007/BF01208923. [[CrossRef](#)] [[Google Scholar](#)]
128. Luts A., Salm J. Chemical composition of small atmospheric ions near the ground. *J. Geophys. Res.* 1994;99:10781–10785. doi: 10.1029/93JD03225. [[CrossRef](#)] [[Google Scholar](#)]
129. Goldstein N., Lewin T., Kamensky A., Dubinin V., Baumann S., Konstantinova O. Exogenous gaseous superoxide potentiates the antinociceptive effect of opioid analgesic agents. *Inflamm. Res.* 1996;45:473–478. doi: 10.1007/BF02252319. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
130. Byczkowski J.Z., Gessner T. Biological role of superoxide ion-radical. *Int. J. Biochem.* 1988;20:569–580. doi: 10.1016/0020-711X(88)90095-X. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
131. Fridovich I. Superoxide dismutases. An adaptation to a paramagnetic gas. *J. Biol. Chem.* 1989;264:7761–7764. [[PubMed](#)] [[Google Scholar](#)]
132. Lim S.S., Vos T., Flaxman A.D., Danaei G., Shibuya K., Adair-Rohani H., Amann M., Anderson H.R., Andrews K.G., Aryee M., et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet.* 2012;380:2224–2260. doi: 10.1016/S0140-6736(12)61766-8. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

133. Chow J.C., Watson J.G., Mauderly J.L., Costa D.L., Wyzga R.E., Vedal S., Hidy G.M., Altshuler S.L., Marrack D., Heuss J.M., et al. Health effects of fine particulate air pollution: Lines that connect. *J. Air Waste Manage. Assoc.* 2006;56:1368–1380. doi: 10.1080/10473289.2006.10464545. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
134. Xie Y., Zhao B., Zhang L., Luo R. Spatiotemporal variations of PM_{2.5} and PM₁₀ concentrations between 31 Chinese cities and their relationships with SO₂, NO₂, CO and O₃. *Particuology*. 2015;20:141–149. doi: 10.1016/j.partic.2015.01.003. [[CrossRef](#)] [[Google Scholar](#)]
135. Xing Y.F., Xu Y.H., Shi M.H., Lian Y.X. The impact of PM_{2.5} on the human respiratory system. *J. Thorac. Dis.* 2016;8:E69–E74. doi: 10.3978/j.issn.2072-1439.2016.01.19. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
136. Wang N., Mengersen K., Kimlin M., Zhou M., Tong S., Fang L., Wang B., Hu W. Lung cancer and particulate pollution: A critical review of spatial and temporal analysis evidence. *Environ. Res.* 2018;164:585–596. doi: 10.1016/j.envres.2018.03.034. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
137. Erqou S., Clougherty J.E., Olafiranye O., Magnani J.W., Aiyer A., Tripathy S., Kinnee E., Kip K.E., Reis S.E. Particulate matter air pollution and racial differences in cardiovascular disease risk. *Arterioscler. Thromb. Vasc. Biol.* 2018;38:935–942. doi: 10.1161/ATVBAHA.117.310305. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
138. Lee B.U., Yermakov M., Grinshpun S.A. Removal of fine and ultrafine particles from indoor air environments by the unipolar ion emission. *Atmos Environ.* 2004;38:4815–4823. doi: 10.1016/j.atmosenv.2004.06.010. [[CrossRef](#)] [[Google Scholar](#)]
139. Mayya Y.S., Sapra B.K., Khan A., Sunny F. Aerosol removal by unipolar ionization in indoor environments. *Aerosol Sci.* 2004;35:923–941. doi: 10.1016/j.jaerosci.2004.03.001. [[CrossRef](#)] [[Google Scholar](#)]
140. Grinshpun S.A., Mainelis G., Trunov M., Adhikari A., Reponen T., Willeke K. Evaluation of ionic air purifiers for reducing aerosol exposure in confined indoor spaces. *Indoor Air.* 2005;15:235–245. doi: 10.1111/j.1600-0668.2005.00364.x. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
141. Berry D., Mainelis G., Fennell D. Effect of an ionic air cleaner on indoor/outdoor particle ratios in a residential environment. *Aerosol Sci. Technol.* 2007;41:315–328. doi: 10.1080/02786820701199702. [[CrossRef](#)] [[Google Scholar](#)]
142. Qiu B., Li Q., Hong W., Xing G. Characterization of the key material for elimination of PM_{2.5} particles in the atmosphere. *J. Spectrosc.* 2015 doi: 10.1155/2015/472019. [[CrossRef](#)] [[Google Scholar](#)]
143. Tanaka A., Zhang Y. Dust settling efficiency and electrostatic effect of a negative ionization system. *J. Agr. Saf. Health.* 1996;2:39–47. doi: 10.13031/2013.19440. [[CrossRef](#)] [[Google Scholar](#)]
144. Grabarczyk Z. Electiveness of indoor air cleaning with corona ionizers. *J. Electrostat.* 2001;51:278–283. doi: 10.1016/S0304-3886(01)00058-4. [[CrossRef](#)] [[Google Scholar](#)]
145. Niu J.L., Tung T.C.W., Burnett J. Quantification of dust removal and ozone emission of ionizer air-cleaners by chamber testing. *J. Electrostatics.* 2001;51:20–24. doi: 10.1016/S0304-3886(01)00118-8. [[CrossRef](#)] [[Google Scholar](#)]
146. Daniels S.L. On the ionization of air for removal of noxious effluvia (Air ionization of indoor environments for control of volatile and particulate contaminants with nonthermal plasmas generated by dielectric-barrier discharge. *IEEE T. Plasma Sci.* 2002;30:1471–1481. doi: 10.1109/TPS.2002.804211. [[CrossRef](#)] [[Google Scholar](#)]

147. Sawant V.S., Meena G.S., Jadhav D.B. Effect of negative air ions on fog and smoke. *Aerosol Air Qual. Res.* 2012;12:1007–1015. doi: 10.4209/aaqr.2011.11.0214. [[CrossRef](#)] [[Google Scholar](#)]
148. Sawant V.S. Control of respirable particles in indoor air with portable negative air ion generator. *IOSR J. Environ. Sci. Toxicol. Food Technol.* 2013;3:28–31. doi: 10.9790/2402-0332831. [[CrossRef](#)] [[Google Scholar](#)]
149. Sawant V.S. Removal of particulate matter by using negative electric discharge. *Int. J. Eng. Innov. Technol.* 2013;2:48–51. [[Google Scholar](#)]
150. Černecký J., Valentová K., Pivarčiová E., Božek P. Ionization impact on the air cleaning efficiency in the interior. *Meas. Sci. Rev.* 2015;15:156–166. doi: 10.1515/msr-2015-0023. [[CrossRef](#)] [[Google Scholar](#)]
151. Pushpawela B., Jayaratne R., Nguy A., Morawska L. Efficiency of ionizers in removing airborne particles in indoor environments. *J. Electrostatics.* 2017;90:79–84. doi: 10.1016/j.elstat.2017.10.002. [[CrossRef](#)] [[Google Scholar](#)]
152. Zhao X., Li Y., Hua T., Jiang P., Yin X., Yu J., Ding B. Low-resistance dual-purpose air filter releasing negative ions and effectively capturing PM_{2.5}. *ACS Appl. Mater. Interfaces.* 2017;9:12054–12063. doi: 10.1021/acsami.7b00351. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
153. Yu K.P., Lee W.M.G., Peng C.J., Chen Y.C., Shen W.T. Effects of roughness, dielectric constant and electrical resistivity of wall on deposition of submicron particles driven by ionic air purifier. *J. Environ. Chem. Engineer.* 2017;5:3108–3114. doi: 10.1016/j.jece.2017.06.024. [[CrossRef](#)] [[Google Scholar](#)]
154. Sutto H.C. Superoxide and ozone production by corona discharge. *J. Chem. Soc. Faraday Trans.* 1984;80:2301–2311. doi: 10.1039/f19848002301. [[CrossRef](#)] [[Google Scholar](#)]
155. Wu C.C., Lee G.W.M. Oxidation of Volatile Organic Compounds by Negative Air Ions. *Atmos. Environ.* 2004;38:6287–6295. doi: 10.1016/j.atmosenv.2004.08.035. [[CrossRef](#)] [[Google Scholar](#)]
156. Waring M.S., Siegel J.A., Corsi R.L. Ultrafine particle removal and generation by portable air cleaners. *Atmos. Environ.* 2008;42:5003–5014. doi: 10.1016/j.atmosenv.2008.02.011. [[CrossRef](#)] [[Google Scholar](#)]
157. Yu K.P. Enhancement of the deposition of ultrafine secondary organic aerosols by the negative air ion and the effect of relative humidity. *J. Air Waste Manage.* 2012;62:1296–1304. doi: 10.1080/10962247.2012.708006. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]